



Europäisches Patentamt  
European Patent Office  
Office européen des brevets

(11) Publication number:

0 209 234  
A2

1

(12)

## EUROPEAN PATENT APPLICATION

(21) Application number: 86304263.6

(22) Date of filing: 04.06.86

(51) Int. Cl.: A 01 N 47/44, A 01 N 47/26,  
A 01 N 43/78  
//  
(A01N47/44, 43:78), (A01N47/26,  
43:78), (A01N43/78, 43:653,  
43:647, 43:54, 43:52, 43:50, 43:36,  
37:30, 37:04, 33:18)

(30) Priority: 05.06.85 GB 8514217  
30.04.86 GB 8610641

(43) Date of publication of application: 21.01.87  
Bulletin 87/4

(54) Designated Contracting States: DE FR GB IT

(71) Applicant: UNIROYAL LIMITED, Cliftonhall Road,  
Newbridge Midlothian EH28 8TW Scotland (GB)

(72) Inventor: Bartlett, Digby H., 1 Hailsham Close,  
Battenhall Worcester WR8 2JB (GB)

(74) Representative: Harrison, Michael Robert et al,  
Urquhart-Dykes & Lord 5th Floor Tower House Merlion  
Way, Leeds, LS2 8PA (GB)

(54) Synergistic fungicidal composition.

(57) Compositions comprised of (A) certain substituted carbox-amidothiazoles and (B) one or more compounds selected from the group consisting of imidazoles, triazoles, guanadines, halonitrobenzenes, manganese and zinc salts of ethylene-bis-(dithiocarbamate), pyrimidines, dicarboxamides, pyrrolidines and thiurams exhibit synergistic fungicidal activity. Also, a method for protecting plants from fungal disease comprising treating such plants with a fungicidally effective amount of said composition is disclosed.

EP 0 209 234 A2

ACTORUM AG

SYNERGISTIC FUNGICIDAL COMPOSITIONFIELD OF THE INVENTION

5 This invention relates to a synergistic fungicidal composition comprised of (A) certain substituted carbox-amidothiazoles and (B) one or more compounds selected from the group consisting of imidazoles, triazoles, 10 guanadines, halonitrobenzenes, manganese and zinc salts of ethylene-bis(dithiocarbamate), pyrimidines, dicarbox-amides, pyrrolidines and thiurams. In another aspect, this invention relates to a method for protecting plants from fungal disease, which method involves treating such 15 plants with a fungicidally effective amount of said composition.

BACKGROUND OF THE INVENTION

20 The need for effective protection of plants against fungal disease requires no amplification, the results of such fungal attack being well documented. However, in the past, many effective fungicides were mercury-based, and were thus toxic to warm-blooded animals.

25 Accordingly, several non-mercury-based fungicides have been developed. Thus, for example, U.S. Patent 3,505,055 to von Schmeling et al discloses carboxamido-thiazoles which are effective fungicides. Similarly,


U.S. Patent 3,658,813 to Godefroi et al shows fungicidal  
1-[beta-aryl-beta-(R-oxy)-ethyl] imidazoles, while  
British Patent 1,522,657 discloses dioxolan-substituted  
triazoles which exhibit fungicidal activity. Sarrett et  
al, in U.S. Patent 3,017,415, show certain thiazolyl  
benzimidazoles which function as anthelmintics.

While the above compounds will function admirably as  
fungicides when employed alone, it has been unexpectedly  
found that when certain compounds of the latter three  
patents (along with certain other known compounds) are  
employed in conjunction with certain carboxamidothiazoles  
(some of which are disclosed in U.S. Patent 3,505,055 and  
others of which are closely related compounds) the  
efficacy of such combinations is greater than that  
predicted for the mere addition of the compounds alone.

#### DESCRIPTION OF THE INVENTION

In one aspect, the present invention is directed to  
a fungicidal composition comprising:

(A) a compound of the formula:




wherein R and R<sup>1</sup> are the same or different and  
are hydrogen or C<sub>1</sub>-C<sub>8</sub> alkyl; and

- (B) one or more compounds selected from the group consisting of imidazoles, triazoles, guanadines, halonitrobenzenes, manganese and zinc salts of ethylene-bis(dithiocarbamate), pyrimidines, dicarboxamides, pyrrolidines and thiurams.

In another aspect, this invention relates to a method for protecting plants from fungal disease, which method comprises applying to such plants a fungicidally effective amount of a composition comprising:

- (A) a compound of the formula:




wherein R and R<sup>1</sup> are the same or different and are hydrogen or C<sub>1</sub>-C<sub>8</sub> alkyl; and

- (B) one or more compounds selected from the group consisting of imidazoles, triazoles, guanadines, halonitrobenzenes, manganese and zinc salts of ethylene-bis(dithiocarbamate), pyrimidines, dicarboxamides, pyrrolidines and thiurams.

The carboxamidothiazoles which may be employed in the present invention are disclosed in U.S. Patent 3,505,055 or are closely related thereto and are compounds of the formula:

0209234



5 wherein R and R<sup>1</sup> may be the same or different and are hydrogen or C<sub>1</sub>-C<sub>8</sub> alkyl.

More preferably, R and R<sup>1</sup> are each independently hydrogen or C<sub>1</sub>-C<sub>4</sub> alkyl. Most preferably, R and R<sup>1</sup> are both methyl.

The imidazoles, triazoles, guanadines, halonitrobenzenes, manganese and zinc salts of ethylenebis(dithiocarbamate), pyrimidines, dicarboxamides, or thiurams which comprise component (B) of the composition of the present invention are all known compounds having fungicidal activity.

Representative of the imidazoles which may be employed are 1-[2-(2,4-dichlorophenyl)-2-(2-propenyloxy)ethyl]-1H-imidazole, 1-N-propyl-N-[2-(2,4,6-trichlorophenoxy)ethyl] carbamoylimidazole, 1-[(2-[4-(4-bromophenoxy)phenyl]-1,3-oxathiolan-2-yl)methyl]-1H-imidazole, 1-[(2-[4-fluorophenyl]-1,3-oxathiolan-2-yl)methyl]-1H-imidazole, and 2-(4-thiazolyl)-1H-benzimidazole. Illustrative of the triazoles which may be employed are 1-[[2-(2,4-dichlorophenyl)-4-ethyl-1,3-dioxolan-2-yl]methyl]-1H-1,2,4-triazole, 1-[[2-(1,1'-biphenyl-4-yl)-1,3-oxathiolan-2-yl]methyl]-1H-1,2,4-

triazole, 1-[[2-(4-fluorophenyl)-1,3-oxathiolan-2-yl]methyl]-1H-1,2,4-triazole, 1-[[2-(4-phenoxyphenyl)-1,3-dithiolan-2-yl]methyl]-1H-1,2,4-triazole, 1-[(2-[[[(1,1'-biphenyl)-4-yl]-1,3-dithian-2-yl]methyl]-1H-1,2,4-triazole, 1-[[2-[2-(1-methoxy)phenyl]-1,3-oxathiolan-2-yl]methyl]-1H-1,2,4-triazole, 4-chlorobenzyl N-(2,4-dichlorophenyl)-2-(1,2,4-triazol-1-yl)ethanethioimide and 2,4'-difluoro-alpha-(1H-1,2,4-triazole-1-ylmethyl) benzhydrol alcohol.

10 Representative of the guanadines which may be employed is 1,1'-iminobis(octamethylene) diguanadine, and illustrative of the halonitrobenzenes which may be employed is pentachloronitrobenzene.

Component (B) of the composition of this invention  
15 may comprise manganese, zinc or mixed manganese and zinc salts of ethylene-bis(dithiocarbamate).

Illustrative of the pyrimidines which may be employed are 5-butyl-2-ethylamino-6-methyl-4-(3H)-pyrimidinone and alpha-(2-chlorophenyl)-alpha-(4-fluorophenyl)-5-pyrimidine methanol. Representative of the  
20 dicarboxamides which may be employed is 3-(3,5-dichlorophenyl)-N-(1-methylethyl)-2,4-dioxo-1-imidazolidine carboxamide, while tetramethylthiuram is representative of the thiurams which may be employed.

25 Illustrative of the pyrrolidines which may be employed is 1-(3,5-dichlorophenyl)-3-methoxymethyl-2,5-pyrrolidindione.

Generally, the weight ratio of component (A) (i.e., the carboxamidothiazole) to component (B) present in the composition of this invention is preferably between about 1:5 and about 500:1. Typically, said weight ratio is  
- 5 generally more preferably between about 1:2 and about 200:1.

When component (B) is 1-[2-(2,4-dichlorophenyl)-2-(2-propenyloxy)ethyl]-1H-imidazole or 2-(4-thiazolyl)-1H-benzimidazole most preferably the weight ratio of  
10 component (A) to component (B) is between about 5:1 and about 75:1. When component (B) is 1-[[2-(2,4-dichlorophenyl)-4-ethyl-1,3-dioxolan-2-yl]methyl]-1H-1,2,4-triazole, most preferably the weight ratio of component (A) to component (B) is between about 25:1 and about  
15 375:1. When component (B) is 4-chlorobenzyl N-(2,4-dichlorophenyl)-2-(1,2,4-triazol-1-yl)ethanethioimide, most preferably the weight ratio of component (A) to component (B) is between about 1:2 and 10:1. The most preferred weight ratios for any particular set of  
20 components (A) and (B) may be readily determined by one skilled in the art utilizing conventional experimentation.

The composition of the present invention may further comprise one or more carrier materials such as talc,  
25 kaolin, bentonite and baryters, as well as stickers such as Spindeloil, ethylene glycol and polyvinylacetate. The composition may also include conventionally employed

pigments such as iron oxide, Rubin Toner, titanium dioxide, and the like.

The compositions of this invention may be formulated as a powder by mixing together all the dry materials, including the carrier and additives until a homogenous mixture is formed. The stickers are then added and the whole mixed again until the mixture becomes essentially homogenous.

In the case of a liquid formulation, organic solvents such as xylene, dimethyl formamide, ethylene glycol, methanol and cyclohexane can be used as diluting agents and mixed together with surface active agents (such as, for example, calcium dodecylbenzenesulphonate, polyglycol ether and acrylopolyether alcohols) as well as a dye (such as Rhodamine, methylviolet and the like).

Moreover, the composition of this invention may be used together with one or more additional pesticidal materials such as insecticides, for instance organochlorine compounds such as Lindane and the like; organophosphoric esters such as Diazinon, Isazofos, Thiofanox and the like; and carbamates such as Carbofuran, Mercaptodimethar, Bendiocarb and the like; and repellants such as anthraquinone, thioanthraquinone, Benzathrone, Ziram and diphenylguanidine and the like in order to protect the plants against pests such as insects.

The method of this invention comprises applying to plants a fungicidally effective amount of the composition



of this invention. Although the composition may be applied to any portion of the plants to be protected, the instant compositions are especially effective when employed as seed dressings.

5        When employed as seed dressings, preferably between about 5 grams and about 100 grams of composition is applied to each 100 kilograms of seed. More preferably, the composition is applied at a rate of between about 10 grams and about 50 grams per 100 kilograms of seed.

10        It is to be understood that the synergistic fungicidal control exhibited by the compositions of this invention is only exhibited wherein limited amounts of fungicide is employed -- i.e., it is impossible to have synergy wherein the fungicides comprising this composition are used in such large amounts that essentially  
15        complete control would be expected employing either component alone. The essence of this invention is that lesser amounts of fungicide are necessary to obtain a given degree of control.

20        As is demonstrated in the examples below, the compositions of this invention exhibit unexpectedly enhanced, i.e., synergistic activity, rather than merely cumulative effect.

      The compositions and method of this invention are  
25        useful in controlling a wide spectrum of fungi, such as Calonectria nivalis, Pyrenophora graminea, Tilletia caries, Ustilago avenae Septoria nodorum, Tilletia foetida, Ustilago hordei, Ustilago nigra, Urocystis

0209234

occulata, Ustilago nuda, Ustilago tritici, Typhula incarnata, Rhynchosporium secalis, Erysipha graminis and the like, and demonstrate a high degree of activity against any certain seed pathogens, such as Pyrenophora avenae, Cochliobolus savitus and the like.

Moreover, the compositions of this invention are admirably suited for agricultural, horticultural and forestry applications because they are far less toxic to warm-blooded animals than mercury based fungicides.

#### EXAMPLES

The following Example are intended to further illustrate the invention and are not intended to limit the scope of the invention in any manner whatsoever.

#### Examples 1-9

In these Examples, the seed treatment method was carried out in accordance with the BBA guidelines 4-1.1 (March 1974) Ehle, H., Frohbeyer, P.E., Reinhard, H. and Roder, K. (1974), "Preliminary guidelines for the testing of seed dressings against cereal diseases", March 1974, published by BBA, Messeweg 11/12, 3300 Braunschweig. In this example, the following abbreviations represent the below listed compounds:

DMCI = 2,4-dimethyl-5-carboxanilidothiazole  
DCPI = 1-[2-(2,4-dichlorophenyl)-2-(2-propenyloxy)ethyl]-1H-imidazole  
TBIA = 2-(4-thiazolyl)-1H-benzimidazole  
DCDT = 1-[[2-(2,4-dichlorophenyl)-4-ethyl-1,3-dioxolan-2-yl]methyl]-1H-1,2,4-triazole

0209234

A wettable powder formulation was prepared using the following ingredients (all parts by weight):

	DMCI	75.0
5	Surfactants (*)	4.1
	Rhodamine dye 3500	0.5
	Ethylene glycol	3.0
	Dixie Clay	17.4

10 (\*) Mixture of alkali metal salts of alkyl naphthalene sulfonic acids: poly(ethylene oxide - propylene oxide).

Additional formulations were made up by dissolving the component (B) fungicides in dimethylformamide at the various concentrations (percent weight) indicated below (each formulation contained also 0.8% Rhodamine dye 3500):

	<u>Fungicide</u>	<u>Concentrations</u>
	DCPI	0.5
20	TBIA	0.5
	DCDT	0.1

Spring wheat cv. Timmo was artificially infected with dry spores of Tilletia caries at a rate of 2 grams of spores per kilogram of wheat seed. The infected wheat seed was treated with various amounts each of the formulations above as summarized in Table I below. This treatment comprised diluting the above formulations with

water and spraying a fine mist of the particular diluted  
formulation employed on the infected seeds using a  
deVilbiss [trademark] compressed air powered spray gun,  
followed by shaking the seed and chemical in a bottle for  
5 5 minutes.

The seeds, which had been infected and treated as  
described above, were drilled into open ground, on a clay  
loam soil type, using a tractor mounted Oyjord Seed  
Drill. Plot size was 1.6 meters x 5 meters with four  
10 replications, all treatments being fully randomized.

Assessment of fungicidal effect was carried out by  
taking 100 randomly selected heads per plot and examining  
for disease thus establishing the percent control of  
infection. The results of such testing are summarized in  
15 Table I, below.

20

25

0209234

TABLE I

<u>Experiment</u>	<u>A</u>	<u>B</u>	<u>C</u>	<u>D</u>	<u>E</u>	<u>F</u>	<u>G</u>	<u>H</u>	<u>I</u>
DMCI	37.5	75	-	-	-	-	-	-	-
DCPI	-	-	1	5	-	-	-	-	-
TBIA	-	-	-	-	1	3	5	-	-
DCDT	-	-	-	-	-	-	-	0.2	0.6
Actual Control (Percentage)	59.5	95.2	2.4	7.1	0	2.4	38.1	71.4	92.9

Employing the formulations and methods of treatment described in Experiments A - I above, several Experiments were conducted employing a mixture of compounds as summarized in Table II below. The expected control percentages were calculated for each Experiment employing the results of Experiments A - I utilizing the formula developed by S. R. Colby, "Calculating Synergistic and Antagonistic Responses of Herbicide Combinations", Weeds, Vol. 15, pp. 20-22 (1967), the disclosure of which is herein incorporated by reference.

The Colby formula may be summarized as follows:

If X = the percent inhibition of growth by herbicide

A at p lb/A

and Y = the percent inhibition of growth by

herbicide

B at q lb/A

and E = the expected percent inhibition of growth by

herbicides

A + B at p + q lb/A

then,

$$E = X + Y - \frac{XY}{100}$$

When the observed response is greater than expected, the combination is synergistic; when less than expected, it is antagonistic. If the observed and expected responses are equal, the combination is additive.

The expected and actual control percentages for each Experiment are listed in Table II.

0209234

TABLE II

<u>Example No.</u>		<u>1</u>	<u>2</u>	<u>3</u>	<u>4</u>	<u>5</u>	<u>6</u>
5	DMCI	37.5	37.5	75	75	37.5	37.5
	DCPI	1	5	1	5	-	-
	TBIA	-	-	-	-	1	3
	DCDT	-	-	-	-	-	-
	Control:						
10	Expected Percentage	60.5	62.4	95.3	95.5	59.5	60.5
	Actual Percentage	69	85.7	100	97.6	81	71.4
<u>Example No.</u>		<u>7</u>	<u>8</u>	<u>9</u>			
15	DCMI	37.5	37.5	37.5			
	DCPI	-	-	-			
	TBIA	5	-	-			
	DCDT	-	0.2	0.6			
20	Control:						
	Expected Percentage	74.9	88.4	97.1			
	Actual Percentage	92.9	100	100			

The data on Table II is clearly indicative of the synergistic fungicidal effects demonstrated by the compositions of the present invention wherein component (B) is an imidazole or a triazole.

Examples 10-15

The efficacy of several of the compositions of this invention in controlling Fusarium nivale was determined by the following process. Wheat seeds, infected with

- 5 F. nivale were treated with the chemicals in the dosages listed in Table III in a seed treating machine. One hundred seeds for each treatment were placed between two sheets of water-soaked absorbent paper to provide moisture for seed germination. The blotter paper was
- 10 then rolled loosely holding the seeds in the moist paper. The rolled blotters were then placed in a sealable plastic bag with 5 cm space left open at the top for proper air exchange. All the treatments were placed in separate bags in a cold chamber at 5°C for 7 days.
15. Untreated control bags were also done in the same manner. This temperature promoted development of the Fusarium disease. The bags were then removed and transferred to an illuminated incubator at 20°C for another 7 days for further disease development. At the end of this period
- 20 efficacy of control was determined by counting the numbers of infected seedlings from each treatment and tabulating the % diseased seedlings. These data were compared to that from untreated controls and the percentage reduction in disease (% control) was
- 25 calculated for each treatment.

Moreover, the theoretical control for each combination was determined employing the Colby formula. The results of such testing are summarized in Table III.



TABLE III

EXAMPLE	COMPOUND	CONCENTRATION (g/100 kg seed)	% CONTROL	THEORETICAL CONTROL	OBSERVED CONTROL
10A	DMCI <sup>1</sup>	18	30.8	--	--
10B	IODG <sup>2</sup>	30	66.1	--	--
10	DMCI	(18 g/100 kg) + IODG(30 g/100 kg)		76.54	86.1
11A	DMCI	18	30.8	--	--
11B	PCNB <sup>3</sup>	20	16.9	--	--
11	DMCI	(18 g/100 kg) + PCNB(20 g/100 kg)		42.5	50.8
12A	DMCI	18	30.8	--	--
12B	MNB <sup>4</sup>	40	10.8	--	--
12	DMCI	(18g/100 kg) + MNB(40g/100 kg)		38.3	50.8
13A	DMCI	18	30.8	--	--
13B	IPRO <sup>5</sup>	15	67.7	--	--
13	DMCI	(18g/100 kg) + IPRO(15g/100 kg)		77.6	80.0

TABLE III (Cont.)

EXAMPLE	COMPOUND	CONCENTRATION (g/100 kg seed)	% CONTROL	THEORETICAL CONTROL	OBSERVED CONTROL
14A	DMCI	18	30.8	--	--
14B	PROC <sup>6</sup>	10	90.8	--	--
14	DMCI (18g/100 kg) + PROC(10g/100 kg)			93.6	100
15A	DMCI	20	71.1	--	--
15B	CBDT <sup>7</sup>	10	0	--	--
15	DMCI (20g/100 kg) + CBDT(10g/100 kg)			71.1	86.5
1	DMCI = 2,4-dimethyl-5-carboxanilidothiazole				
2	IODG = 1,1'-iminobis(octamethylene) diguanadine				
3	PCMB = pentachloronitrobenzene				
4	MNB = manganese salt of ethylene-bis(dithiocarbamate)				
5	I PRO = 3-(3,5-dichlorophenyl)-N-(1-methylethyl)-2,4-dioxo- 1-imidazolidine carboxamide				
6	PROC = 1-N-propyl-N-[2-(2,4,6-trichlorophenoxy)ethyl]carbamoyl- imidazole				
7	CBDT = 4-chlorobenzyl N-(2,4-dichlorophenyl)-2-(1,2,4-triazol- 1-yl)ethanethioimide				

Examples 16-19

Employing the Fusarium nivale test employed in Examples 10-17, several combinations of 2,4-dimethyl-5-carboxanilidothiazole (DMCI) and 4-chlorobenzyl N-(2,4-dichlorophenyl)-2-(1,2,4-triazol-1-yl)ethanethioimide (CBDT) were evaluated for their fungicidal efficacy. The results of such testing are summarized in Table IV.

The control observed for DMCI alone is as follows:

10	<u>Grams DMCI/100 kg seed</u>	<u>% control</u>
	20	71.1
	40	94.2

The control observed for CBDT alone is as follows:

15	<u>Grams CBDT/100 kg seed</u>	<u>% control</u>
	10	0
	20	28.8

20

25

TABLE IV

<u>Example</u>	<u>Grams DMCI per 100kg seed</u>	<u>Grams CBDT per 100kg seed</u>	<u>Theoretical % Control</u>	<u>Observed % Control</u>
5	16	20	71.10	86.5
	17	20	79.42	94.2
	18	40	94.20	100
	19	40	95.88	100

10        The above results indicate the synergistic effect of  
a composition comprising DMCI and CBDT.

         The above data indicate the unexpectedly superior  
control exhibited by the compositions of this invention  
relative to the control exhibited by either of the  
15 components alone.

20

25


## CLAIMS:

1. A fungicidal composition characterised in that it comprises:

(A) a compound of the formula:

5

10



15

wherein R and R¹ are the same or different and are  
20 hydrogen or C₁-C₈ alkyl; and

(B) one or more compounds selected from imidazoles, triazoles, guanadines, halonitrobenzenes, manganese and zinc salts of ethylene-bis(dithiocarbamate), pyrimidines, dicarboxamides, pyrrolidines and thiurams.

25 2. A composition according to claim 1 characterised in that R and R¹ are hydrogen or C₁-C₄ alkyl.

3. A composition according to claim 1 or claim 2 characterised in that R and R¹ are methyl.

4. A composition according to any of the preceding  
30 claims characterised in that the weight ratio of compound (a) to compound (b) is between about 1:5 and about 500:1.

5. A composition according to any of the preceding claims characterised in that the weight ratio of compound (a) to compound (b) is between about 1:2 and 200:1.

35 6. A fungicidal composition according to any of the

preceding claims characterised in that said imidazole is selected from 1-[2,4-dichlorophenyl)-2-(2-propenyloxy)ethyl]-1H-imidazole, 1-N-propyl-N-[2-(2,4,6-trichlorophenoxy)ethyl] carbamoylimidazole, 1-[(2-[4-(4-bromophenoxy)phenyl-1,3-oxathiolan-2-yl)methyl]-1H-imidazole, 1-[(2-[4-fluorophenyl]-1,3-oxathiolan-2-yl)methyl]-1H-imidazole, 2-(4-thiazolyl)-1H-benzimidazole, 1-[[2-(2,4-dichlorophenyl)-4-ethyl-1,3-dioxolan-2-yl)methyl]-1H-1,2,4-triazole, 1-[[2-(1,1'-biphenyl-4-yl)-1,3-oxathiolan-2-yl)methyl]-1H-1,2,4-triazole, 1-[[2-(4-fluorophenyl)-1,3-oxathiolan-2-yl)methyl]-1H-1,2,4-triazole, 1-[[2-(4-phenoxyphenyl)-1,3-dithiolan-2-yl)methyl]-1H-1,2,4-triazole, 1-[(2-[[1,1'-biphenyl]-4-yl]-1,3-dithian-2-yl)methyl]-1H-1,2,4-triazole, 1-[[2-[2-(1-methyl ethoxy)phenyl]-1,3-oxathiolan-2-yl)methyl]-1H-1,2,4-triazole, 4-chlorobenzyl N-(2,4-dichlorophenyl)-2-(1,2,4-triazol-1-yl) ethanethioimide, 2,4'-difluoro-alpha-(1H-1,2,4-triazole-1-ylmethyl)benzhydryl alcohol, 1,1'-iminobis(octamethylene) diguanadine, pentachloronitrobenzene, a manganese, zinc or mixed manganese and zinc salt of ethylene-bis(dithiocarbamate), 5-butyl-2-ethylamino-6-methyl-4-(3H)-pyrimidinone, alpha-(2-chlorophenyl)-alpha-(4-fluorophenyl)-5-pyrimidine methanol, 3-(3,5-dichlorophenyl)-N-(1-methyl-ethyl)-2,4-dioxo-1-imidazolidine carboxamide, tetramethylthiuram, and 1-(3,5-dichlorophenyl)-1-methoxy-methyl-2,4-pyrrolidindione.

7. A method for protecting plants from fungal disease, characterised in that the method comprises applying to such plants a fungicidally effective amount of a composition as claimed in any of the preceding claims.



Europäisches Patentamt  
European Patent Office  
Office européen des brevets

(11) Publication number:

**0 209 234  
A3**

(12) **EUROPEAN PATENT APPLICATION**

(21) Application number: **86304263.6**

(22) Date of filing: **04.06.86**

(51) Int. Cl.<sup>4</sup>: **A 01 N 47/44**  
**A 01 N 47/26, A 01 N 43/78**  
**/(A01N47/44, 43:78),**  
**(A01N47/26, 43:78), (A01N43/78,**  
**43:653, 43:647, 43:54, 43:52,**  
**43:50, 43:36, 37:30, 37:04,**  
**33:18)**

(30) Priority: **05.06.85 GB 8514217**  
**30.04.86 GB 8610541**

(43) Date of publication of application:  
**21.01.87 Bulletin 87/4**

(88) Date of deferred publication of search report: **20.05.87**

(84) Designated Contracting States:  
**DE FR GB IT**

(71) Applicant: **UNIROYAL LIMITED**  
**Cliftonhall Road**  
**Newbridge Midlothian EH28 8TW Scotland(GB)**

(72) Inventor: **Bartlett, Digby H.**  
**1 Hailsham Close**  
**Battenhall Worcester WR5 2JB(GB)**

(74) Representative: **Harrison, Michael Robert et al,**  
**Urquhart-Dykes & Lord 5th Floor Tower House Merion**  
**Way**  
**Leeds, LS2 8PA(GB)**

(54) **Synergistic fungicidal composition.**

(57) Compositions comprised of (A) certain substituted carboxamidothiazoles and (B) one or more compounds selected from the group consisting of imidazoles, triazoles, guanadines, halonitrobenzenes, manganese and zinc salts of ethylene-bis(dithiocarbamate), pyrimidines, dicarboxamides, pyrrolidines and thiurams exhibit synergistic fungicidal activity. Also, a method for protecting plants from fungal disease comprising treating such plants with a fungicidally effective amount of said composition is disclosed.



0209234

Application number

EP 86 30 4263

DOCUMENTS CONSIDERED TO BE RELEVANT					
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)		
A	CHEMICAL ABSTRACTS, vol. 83, no. 7, August 18, 1975, page 137, ref.no. 54385m; Columbus, Ohio, US M. STEC et al.: "Comparison of the systemic fungicidal activity of 2-amino-4-methyl-5-carboxyanilido-thiazole (ALF) and 2,4-dimethyl-5-carboxyanilidothiazole (ALG) in green house and field tests" & ACTA PHYTOPATHOL. 1973, 8(3-4), 283-94.  * Abstract *	1-7	A 01 N 47/44 A 01 N 47/26 A 01 N 43/78/ (A 01 N 47/44 43:78) (A 01 N 47/26 43:78) (A 01 N 43/78 43:653 43:647 43:54 43:52 43:50 43:36 37:30 37:04 33:18)		
A	GB-A-1 459 116 (HOECHST)  * Page 1, line 7 - page 3, line 1; claims *	1-3			
D,A	US-A-3 505 055 (B. VON SCHMELING et al.)  * Column 2, lines 16-21; claims 1, 11 *	1-3		TECHNICAL FIELDS SEARCHED (Int. Cl. 4)	
A	EP-A-0 006 012 (UNIROYAL Ltd.)  * Claims *	1-7		A 01 N	
<del>XXXXXXXXXXXXXXXXXXXXXXXXXXXXXXX</del>					
Place of search		Date of completion of the search	Examiner		
The Hague		24-09-1986	FLETCHER		
CATEGORY OF CITED DOCUMENTS					
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document			T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons  & : member of the same patent family, corresponding document		





### CLAIMS INCURRING FEES

The present European patent application comprised at the time of filing more than ten claims.

- ☐ All claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for all claims.
- ☐ Only part of the claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for the first ten claims and for those claims for which claims fees have been paid, namely claims:
- ☐ No claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for the first ten claims.

### X LACK OF UNITY OF INVENTION

The Search Division considers that the present European patent application does not comply with the requirement of unity of invention and relates to several inventions or groups of inventions.

namely: Composition comprising (A) and (B):

- 1) Claims 1-7: ..... imidazoles
- 2) Claims 1-7: ..... triazoles
- 3) Claims 1-7: ..... guanidines
- 4) Claims 1-7: ..... halonitrobenzenes
- 5) Claims 1-7: ..... salts of ethylene-bis(dithiocarbamate)
- 6) Claims 1-7: ..... pyrimidines
- 7) Claims 1-7: ..... dicarboxamides
- 8) Claims 1-7: ..... pyrrolidines
- 9) Claims 1-7: ..... thiurams

- ☐ All further search fees have been paid within the fixed time limit. The present European search report has been drawn up for all claims.
- ☐ Only part of the further search fees have been paid within the fixed time limit. The present European search report has been drawn up for those parts of the European patent application which relate to the inventions in respect of which search fees have been paid, namely claims:
- ☒ None of the further search fees has been paid within the fixed time limit. The present European search report has been drawn up for those parts of the European patent application which relate to the invention first mentioned in the claims.

namely claims: 1-7 (mentioned under point 1))

